



ZAMPHIA

ZAMBIA POPULATION-BASED HIV IMPACT ASSESSMENT 2021



DATA MANUAL
SUPPLEMENT
ADDITIONAL TABLE
SPECIFICATIONS

PARTNERS



Table 12.5: Male circumcision

Percent distribution of men aged 15 years and older by self-reported circumcision status, by result of ZAMPHIA HIV test and selected demographic characteristics, ZAMPHIA 2021

Characteristic	Circumcised ¹		Uncircumcised	Total	Number
	Medical circumcision	Nonmedical circumcision			
Result of ZAMPHIA 2021 HIV test					
HIV positive					
HIV negative					
Not tested					
Age of circumcision					
0-4					
5-9					
10+					
Residence					
Urban					
Rural					
Province					
Central					
Copperbelt					
Eastern					
Lusapula					
Lusaka					
Muchinga					
Northern					
North-Western					
Southern					
Western					
Marital status					
Never married					
Married or living together					
Divorced or separated					
Widowed					
Education					
No education					
Primary					
Secondary					
More than secondary					
Wealth quintile					
Lowest					
Second					
Middle					
Fourth					
Highest					
Age					
15-19					
20-24					
25-29					
30-34					
35-39					
40-44					
45-49					
50-54					
55-59					
60-64					
65+					
Total 15-24					
Total 15-49					
Total 50+					
Total 15+					

¹Relates to Global AIDS Monitoring 2020 indicator 3.16: Prevalence of male circumcision; and PEPFAR indicator VMCM_TOTALCIRC NAT / SUBNAT: Total number of men ever circumcised.

Note: Education categories refer to the highest level of education attended, whether or not that level was completed.

[Return to TOC](#)

Datasets and variables used

Dataset adultind
Subset indstatus = 1 and age ≥ 15 and gender = 1 and mcdetail = 1, 2, 3
Analytic Variables mcdetail

Row stratification variables hivstatusfinal
mcmcdage_za/ mctradage_za
urban
hprovince_za
married
education
wealthquintile
agegroup
age15_24, age15_49, age50_up, age15_up (age)

Column stratification variables

<none>
Weight variables intwt0, intwt001, ..., intwt197

